WVS MILK QUALITY LAB

95 S Harris Ave. Suite 203 Waupun, WI 53963 PH: 920-324-3831 e-mail: milklab@waupunvet.com

Producer Name									Routine culture				
Submitter									Mycoplasma				
Email									Prototheca				
Date Samples									Other				
Sample #	Cow #		Conta	agious	6	Environmenta			als	Miscellaneous			
		Strep ag	Staph aureus	Mycoplasma	C. bovis	Strep sp	Staph sp	Coliforms	T pyogenes	Prototheca	Other	No Growth	Comments
1													
2													
3													
4													
5													
6													
7													
8													
9													
10													
11													
12													
13													
14													
15													
16													
17													
18													
19													
20													
21													
22													
23		1											
24		1											
25		1											

Result Interpretation: +4 Numerous +3 Moderate +2 Few +1 Very Few Contagious: Spread from cow to cow during milking process. Causes high SCC and mastitis in herds. Needs immediate attention via treatment, segregation or culling.

Spread from cow to cow during miking process. Causes high SCC and mastins in herds. Needs immediate attention via treatment, segregation of Consult with your veterinarian. Bacteria from cow housing. Treatment of little benefit. Milking dirty or improperly sanitized teats, poor milking routine, heat and mud holes, dirty housing or poor functioning milking equipment can all contribute to this type of mastitis. Unsterile sample collection, improper cooling of sample, improper storage or handling of samples. These cows should be recultured immediately. Environmental:

Contaminated: